

CERTIFIED TRUE COF

# **Republic of the Philippines** Department of Health **OFFICE OF THE SECRETARY**

NOV 2 5 2020

m

**ADMINISTRATIVE ORDER** No. 2020 - 0013 -B

#### Further Amendment to Administrative Order No. 2020-0013 dated 09 **SUBJECT:** April 2020 entitled, "Revised Administrative Order No. 2020-0012, 'Guidelines for the Implementation for the Inclusion of the Coronavirus Disease 2019 (COVID-19) in the list of Notifiable Diseases for Mandatory Reporting to the Department of Health' dated March 17, 2020."

The Administrative Order No. 2020-0013 dated 09 April 2020 entitled, "Revised Administrative Order No. 2020- 0012, 'Guidelines for the Inclusion of the Coronavirus Disease 2019 (COVID-19) in the list of Notifiable Diseases for Mandatory Reporting to the Department of Health' dated March 17, 2020" is further amended to correct the provisions as follows:

Provisions to be Amended	Amended to
VI. SPECIFIC GUIDELINES	VI. SPECIFIC GUIDELINES
A. COVID-19 Surveillance System	A. COVID-19 Surveillance System
2. Case Definition	2. Case Definition
2.1. Suspect case - is a person who is	2.1. Suspect case
presenting with any of the conditions	a. A person who meets the clinical
below:	AND epidemiological criteria:
a. All SARI cases where NO other	
etiology fully explains the clinical	<u>Clinical criteria:</u>
presentation.	1. Acute onset of fever AND
b. ILI case with any one of the	cough;
following:	OR
ii. with no other etiology that fully	2. Acute onset of ANY THREE
explains the clinical presentation	<b>OR MORE</b> of the following
AND a history of travel to or	signs or symptoms: fever,
residence in an area that reported	cough, general
local transmission of COVID-19	weakness/fatigue, headache,
disease during the 14 days prior to	myalgia, sore throat, coryza,
symptom onset OR	dyspnea,
iii. with contact to a confirmed or	anorexia/nausea/vomiting,
probable case of COVID-19 in the	diarrhea, altered mental status
two days prior to onset of illness of	
the probable/confirmed COVID-19	AND
case became negative on repeat	Epidemiological criteria:
testing.	1. Residing or working in an area
Individuals with fever or cough or	with high risk of transmission
shortness of breath or other	of the virus: for example, closed
respiratory signs or symptoms	residential settings and

Direct Line: 711-9502; 711-9503 Fax: 743-1829 • URL: http://www.doh.gov.ph; e-mail: ftduque@doh.gov.ph

<ul> <li>i. Aged 60 years and above ii. With a comorbidity iii. Assessed as having a high-risk pregnancy iv. Health worker</li> <li>2. Residing in or travel to an area with community transmission anytime within the 14 days prior to symptom onset;</li> <li>OR</li> <li>2. Residing in or travel to an area with community transmission anytime within the 14 days prior to symptom onset;</li> <li>OR</li> <li>3. Working in health setting, including within health facilities and within households, anytime within the 14 days prior to symptom onset;</li> <li>OR</li> <li>3. Working in health setting, including within health facilities and within the 14 days prior to symptom onset.</li> <li>b. A patient with severe acute respiratory illness (SARI: acute support of the stillness illness is based on current availlable data on COVID-19 include the following (Manna 2020):</li></ul>		
<ul> <li>2.2. Probable case - a suspect case who fulfills anyone of the following listed below         <ul> <li>a. Suspect case whom testing for COVID-19 is inconclusive</li> <li>b. Suspect who tested positive for COVID-19 but whose test was not conducted in a national or subnational reference laboratory or officially accredited laboratory for COVID-19 confirmatory testing</li> </ul> </li> <li>CERTIFIED TRUE COPY NOV 277 2020</li> </ul>	conditions: i. Aged 60 years and above ii. With a comorbidity iii. Assessed as having a high pregnancy	<ul> <li>camp and camp-like settings for displaced persons, any time within the 14 days prior to symptom onset;</li> <li>OR</li> <li>2. Residing in or travel to an area with community transmission anytime within the 14 days prior to symptom onset;</li> <li>OR</li> <li>3. Working in health setting, including within health facilities and within households, anytime within the 14 days prior to</li> </ul>
fulfills anyone of the following listed below2.2. Probable casea. Suspect case whom testing for COVID-19 is inconclusivea. A patient who meets clinical criteria above AND is a contact of a probable or confirmed case, or epidemiologically linked to a cluster of cases which has had at least one confirmed case identified within that cluster. Epidemiologically linked refers to exposure of a suspect case to a confirmed case which occurred within 2-14 days prior to the suspect case's onset of illness. This is based on current available data on COVID-19 incubation period.CERTIFIED TRUE COPYNOV 27 2020NOV 27 2020NOV 27 2020		respiratory illness (SARI: acute respiratory infection with history of fever or measured fever of ≥ 38 C°; and cough; with onset within the last 10 days; and who requires hospitalization).
<ul> <li>a. Suspect case whom testing for COVID-19 is inconclusive</li> <li>b. Suspect who tested positive for COVID-19 but whose test was not conducted in a national or subnational reference laboratory or officially accredited laboratory for COVID-19 confirmatory testing</li> <li>a. A patient who meets clinical criteria above AND is a contact of a probable or confirmed case, or epidemiologically linked to a cluster of cases which has had at least one confirmed case identified within that cluster. Epidemiologically linked refers to exposure of a suspect case to a confirmed case which occurred within 2-14 days prior to the suspect case's onset of illness. This is based on current available data on COVID-19 incubation period.</li> <li>b. A suspect case (described above) with chest imaging showing findings suggestive of COVID-19 include the following (Manna 2020):</li> <li>CERTIFIED TRUE COPY</li> <li>NOV 277 2020</li> </ul>	• • • • • • • • • • • • • • • • • • •	
	<ul> <li>a. Suspect case whom testing COVID-19 is inconclusive</li> <li>b. Suspect who tested positive COVID-19 but whose test was conducted in a national subnational reference laboratory officially accredited laboratory COVID-19 confirmatory testing</li> </ul>	a. A patient who meets clinical criteria above AND is a contact of a probable or confirmed case, or epidemiologically linked to a s not or least one confirmed case identified within that cluster.y or v forwithin that cluster. Epidemiologically linked refers to exposure of a suspect case to a confirmed case which occurred within 2-14 days prior to the suspect case's onset of illness. This is based on current available data on COVID-19 incubation period.b. A suspect case (described above) with chest imaging showing findings suggestive of COVID-19 disease**Typical chest imaging findings suggestive of COVID-19 include the following (Manna 2020): • chest radiography: hazy
KMITS - RECORDS SECTION Department of Health	CORAZON'S DELA CRUZ	F mer 2

1

1

<ul> <li>2.3 Confirmed case - any individual, irrespective of presence or absence of clinical signs and symptoms, who was laboratory, and/or DOH-certified laboratory</li> <li>2.3 Confirmed case - any individual, irrespective of presence or absence of clinical signs and symptoms, who was laboratory, and/or DOH-certified laboratory</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of clinical signs and symptoms, who was laboratory, and/or DOH-certified laboratory</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of clinical signs and symptoms, who was laboratory, and/or DOH-certified laboratory.</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of clinical signs and symptoms, who was laboratory, and/or DOH-certified laboratory.</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of clinical signs and symptoms, who was laboratory, and/or DOH-certified laboratory.</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of clinical signs and symptoms, who was laboratory, and/or DOH-certified laboratory.</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of clinical signs and symptoms, who was laboratory, and/or DOH-letensed covel.</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of clinical signs and symptoms.</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of clinical signs and symptoms.</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of aboratory.</li> <li>2.3 Confirmed case + any individual, irrespective of presence or absence of aboratory.</li> <li>3.4 Confirmed case + any individual, irrespective of presence or absence of aboratory.</li> <li>3.5 Confirmed case + any individual, irrespective of presence or absence of aboratory.</li> <li>3.6 Confirmed case + any individual, i</li></ul>	······································	
irrespective of presence or absence of clinical signs and symptoms, who was laboratory confirmed for COVID-19 in a test conducted at the national reference laboratory, a subnational reference laboratory, and/or DOH-certified laboratory testing facility OR any suspect or probable COVID-19 cases, as defined above, who tested positive using antigen tests in areas with outbreaks and/or in remote settings where RT-PCR is not immediately available; provided that the antigen tests satisfy the recommended minimum regulatory, technical and operational specifications set by the Health		<ul> <li>and lower lung distribution</li> <li>chest CT: multiple bilateral ground glass opacities, often rounded in morphology, with peripheral and lower lung distribution</li> <li>lung ultrasound: thickened pleural lines, B lines (multifocal, discrete, or confluent), consolidative patterns with or without air bronchograms.</li> <li>c. A person with recent onset of anosmia (loss of smell) or ageusia (loss of taste) in the absence of any other identified cause.</li> <li>d. Death, not otherwise explained, in an adult with respiratory distress preceding death AND who was a contact of a probable or confirmed case or epidemiologically linked to a cluster which has had at least one confirmed case identified</li> </ul>
any suspect or probable COVID-19 cases, as defined above, who tested positive using antigen tests in areas with outbreaks and/or in remote settings where RT-PCR is not immediately available; provided that the antigen tests satisfy the recommended minimum regulatory, technical and operational specifications set by the Health	irrespective of presence or absence of clinical signs and symptoms, who was laboratory confirmed for COVID-19 in a test conducted at the national reference laboratory, a subnational reference laboratory, and/or DOH-certified laboratory	irrespective of presence or absence of clinical signs and symptoms, who was laboratory confirmed for COVID-19 in a test conducted at the national reference laboratory, a subnational reference laboratory, and/or DOH-licensed
NOV 27 2020	RTIFIED TRUE COPY	any suspect or probable COVID-19 cases, as defined above, who tested positive using antigen tests in areas with outbreaks and/or in remote settings where RT-PCR is not immediately available; provided that the antigen tests satisfy the recommended minimum regulatory, technical and operational

ن ار م

VI. SPECIFIC GUIDELINES A. COVID-19 Surveillance System 2. Case Definition	VI. SPECIFIC GUIDELINES A. COVID-19 Surveillance System 2. Case Definition 2.4. Based on the World Health Organization Clinical Management of COVID-19 dated 27 May 2020, there are four (4) COVID-19 disease severity namely, mild, moderate, severe and critical diseases. As data and knowledge on COVID-19 are fast evolving, definitions of the disease severity will be issued in a separate memorandum.
<ul> <li>VI. SPECIFIC GUIDELINES</li> <li>A. COVID-19 Surveillance System</li> <li>3. Case Detection</li> <li>3.1. SARI and ILI Sites and Other Health Facilities, Providers, and Institution (last paragraph under 3.1) Confirmed COV-19 cases assessed as asymptomatic or clinically recovered</li> </ul>	VI. SPECIFIC GUIDELINES A. COVID-19 Surveillance System 3. Case Detection 3.1. SARI and ILI Sites and Other Health Facilities, Providers, and Institution (last paragraph under 3.1)
by the attending physician shall be tested and will be discharged after at least one negative result. Confirmed COVID-19 case who have clinically recovered or well with negative results on repeat testing shall be reported as RECOVERED. If said discharged cases developed new signs or symptoms or progression from mild to	Criteria for discharge for suspect, probable, and confirmed COVID-19 cases shall no longer entail repeat testing. Criteria for tagging confirmed cases as recovered shall also no longer require repeat testing. <u>Symptomatic patients with mild and</u> <u>moderate symptoms</u> who have:
more serious signs and symptoms, he/she shall be re-admitted once more to isolation and re-testing done. This guideline shall be reviewed and revised accordingly.	a) clinically improved (afebrile for 3 days, clinical improvement of signs and symptoms, and discharged) or are no longer symptomatic for at least three days; AND
CERTIFIED TRUE COPY	b) have completed at least 10 days of isolation either at home, Temporary Treatment and Monitoring Facility (TTMF) or hospital from the onset of illness, can be discharged from isolation and re-integrated to the community without the need for further testing, provided that a
NOV 27 2020	licensed medical doctor clears the patient. Confirmed cases can be
CORAZON S. DELA CRUZ KMITS - RECORDS SECTION Department of Health	to mere 12

do	NR T	
SUE	L S S S S S S S S S S S S S S S S S S S	
F c	S. I. S. S.	
	O O U	
	- Dec Mitts	
CERTIFIED TRUE COPY	CORAZON S. DELA CRUZ KMITS - RECORDS SECTION Department of Health	
	]	

## VI. SPECIFIC GUIDELINES

**C. Recording and Notification System** Health authorities from the government and private sectors, including health facilities, tagged as recovered once these criteria are met.

Symptomatic patients with severe or critical condition who have:

- a) clinically improved (afebrile for 3 days, clinical improvement of signs and symptoms, and discharged) or are no longer symptomatic; AND
- b) have completed at least 21 days of isolation either at home, TTMF or hospital from the onset of illness, can be discharged from isolation and re-integrated to the community without the need for further testing, provided that a licensed medical doctor clears the patient.
- c) Confirmed cases can be tagged as recovered once these criteria are met.

#### **Asymptomatic Patients**

Patients who test PCR positive (+) and remained asymptomatic during the 10 days from date of specimen collection can discontinue isolation after 10 days from date of specimen collection and be tagged as a recovered confirmed case without need for further testing, provided a licensed medical doctor certifies or clears the patient.

Repeat testing among mild and asymptomatic cases should not be a prerequisite for the issuance of a clearance or certification to be issued by medical doctors.

### VI. SPECIFIC GUIDELINES

C. Recording and Notification System

Health authorities from the government and private sectors, including health facilities,

mer 5

laboratory testing facilities, offices, institutions, and individuals, are mandated to report suspect, probable and confirmed cases of COVID-19 and results of COVID-19 testing done within 24 hours of identification or completion of testing.	institutions, and individuals, are mandated to report suspect, probable and confirmed cases of COVID-19 and results of
---	--

• • •

Further, all mentions of COVID-19 Information System pertain to **current information systems** being used by disease reporting units and testing laboratories which include but not limited to COVIDKaya Information System and Laboratory Information System. Further, data management will transition to an EMR-based system with data quality assurance and interoperability to ensure completeness and accuracy of data.

As thus amended, all other provisions stipulated in Administrative Order No. 2020-0013 dated 09 April 2020 and Administrative Order 2020-0013-A date 4 June 2020 not affected by this amendment shall remain in full force and in effect.

This Order shall take effect fifteen (15) days following its publication in the Official Gazette or a newspaper of general circulation.

CISCO/T. DUQUE, III, MD, MSc. Secretary of Health

FIED TRUE COPY NOV 27 CRHZ MITS - RÉCORDS Departme